


FAA Proposed Solar Radiation Alert



Space Weather Week 20 May 2003

Paul Armbruster, ARS-200
Office of Aerospace Weather Policy and Standards



3/6/00

Radiation Exposure of Air Carrier Crewmembers - FAA AC 120-52

RADIATION EXPOSURE OF AIRCREWS AND RECOMMENDED LIMITS

- The recommended annual limit for occupational radiation exposure of an adult is 50 millisieverts (5 rem) (EPA 1987).
- Pregnant crewmembers the dose equivalent to the unborn child from occupational exposure should not be ore than 0.5 millisievert (50 millirem) in any month (NCRP 1987a).

RISK TO CREWMEMBERS

- Death from cancer is the principal health concern associated with occupational exposure to radiation.
- Normal expectation for the U.S. adult population, about 220 of 1000 crew members would die of cancer from causes unrelated to occupational radiation exposure.
- Aircrew members can expect an increased risk of death by cancer on the order of 1 in 360 associated with occupational radiation exposure.

Radiation Exposure of Air Carrier Crewmembers - FAA AC 120-52

- Earth's magnetic field provides some shielding
 - Greatest shielding is over the geomagnetic equator
 - Shielding decreases closer to the poles with little or no change above 50° geographic latitude in North America.
- Earth's atmosphere also provides some shielding
 - Galactic radiation at 1,200 ft. MSL is approximately one-half of one percent of the levels at 39,000 ft. MSL.

Biological Effects

- Unit of measure
 - -1 Sievert (Sv)=100 rem
 - -1 Sv = 1000 mSv
 - $-1 \text{ mSv} = 1000 \mu \text{Sv}$
- Location, Location, Location
 - Altitude, latitude and time dependent
- Solar cycle

Solar Cycle

- 11-year Solar cycle
 - Last maximum occurred Apr 2001
- Radiation from two sources
 - Galactic (cosmic) and Sun (energetic particles)
 - Predominant source dependent on solar cycle
- Cosmic radiation effects are maximum when solar activity is minimum and vice versa

Radiation at Altitude

- Last solar minimum cosmic radiation $4\mu Sv/hr$ at 30K ft at 70N and 3.25 at 35N
- During solar maximum rate fell to about 3μSv/hr at both latitudes
- July 2000 solar particle event 50 μSv/hr extending over almost a full day at altitude

Radiation Limits

- Maximum permissible dose 1000µSv/yr (GP)
- MPD 50,000 μSv/yr for aircrew
- Sea-level exposure of about 6-12µSv/day
- Two-hour flight 20-40K ft 12-24μSv/day
- During solar max it is possible for a solar event to produce a dose rate of up to $200\mu Sv/hr$ for a few hours at airliner altitude

Physiological Impacts

- Cancer leukemia, melanoma, prostate
- Cataracts
- Fetal damage physical, mental, death
- Chromosomal damage Concorde/U2 pilots

Calculating Radiation Exposure

- CAMI-6/6M computer programs:
 - http://jag.cami.jccbi.gov./cariprofile.asp
 - Calculates effective dose of galactic radiation received on an aircraft flying a great circle route (CARI-6) or a user-defined route (CARI-6M)
 - Between any two airports
 - Program accounts for effects of changes in earth's magnetic field and solar activity

Forecasting and Alerts

- CAMI developed plan for alerting on events of at least 20 μ Sv/hr and remaining at or above that level for at least 15 minutes
 - Currently available on:
 - password controlled FTP site for "pull" by airlines
 - FAA Weather Message Switching Center using and SEC supplied WMO message header. (WOXX50)


CAMI Recommended Alert Responses

- Long-lived event (hours)
 - Aircraft in flight descend below 40K ft
 - Aircraft on ground add fuel and change alt
- Short-lived event (minutes)
 - Aircraft in flight descent recommended
 - Aircraft on ground no action

Radiation Exposure of Air Carrier Crewmembers

SOLAR RADIATION ALERT REGIONS

(Alert regions in orange)



Radiation Exposure of Air Carrier Crewmembers - FAA AC 120-52

- •The airline industry agrees that the information on solar radiation is beneficial.
- •The industry is also wary of what, if any, type of regulatory issues may arise from these alerts.
 - •Discussions are underway to resolve these issues.